

Lightning NOx Statistics Derived by NASA Lightning Nitrogen Oxides Model (LNOM) Data Analyses

William Koshak¹ and Harold Peterson²

¹Earth Science Office, ZP11, NASA Marshall Space Flight Center, Robert Cramer Research Hall, 320 Sparkman Drive, Huntsville, AL 35805.

²Universities Space Research Association, Robert Cramer Research Hall, 320 Sparkman Drive, Huntsville, AL 35805.

1. WHAT IS THE LNOM?

The NASA Marshall Space Flight Center (MSFC) Lightning Nitrogen Oxides Model (LNOM) [Koshak et al., 2009, 2010, 2011; Koshak and Peterson 2011, 2013] analyzes VHF Lightning Mapping Array (LMA) and National Lightning Detection Network™ (NLDN) data to estimate the lightning nitrogen oxides (LNOx) produced by individual flashes. **Figure 1** provides an overview of LNOM functionality.




Figure 1. Overview of the LNOM showing input lightning data and output products.

2. BENEFITS OF LNOM

- Does away with unrealistic “vertical stick” lightning channel models for estimating LNOx.
- Uses ground-based VHF data that maps out the true channel in space & time to < 100 m accuracy.
- Therefore, true channel segment height (ambient air density) is used to compute LNOx.
- True channel length is used! (typically tens of kilometers since channel has many branches & “wiggles”).
- Distinction between ground and cloud flashes are made.
- For ground flashes, actual peak current from NLDN used to compute NOx from lightning return stroke.
- NOx computed for several other lightning discharge processes (based on Cooray et al., 2009 theory):
 - Hot core of stepped leaders and dart leaders
 - Corona sheath of stepped leader
 - K-changes
 - Continuing Currents
 - M-components.
- LNOM statistics (see later) can be used to parameterize LNOx production for regional air quality models (like CMAQ), and for global chemical transport models (like GEOS-Chem).

3. STATISTICS

The LNOM data archive at (http://lightning.nsstc.nasa.gov/data/index.html#LNOM_DATA) now contains LNOx analyses for **468,928** flashes. Most of these flashes are derived from 9 years of North Alabama Lightning Mapping Array (Koshak et al, 2004) data as shown below in **Figure 2**; the remaining flashes are from 4 years of data derived from the DC metropolitan LMA network.




Figure 2. LNOM lightning channel length distributions (top 3 plots: all flashes, cloud flashes, ground flashes), and NOx per flash (bottom 3 plots). These distributions and associated statistics are derived from North Alabama thunderstorms.

4. REFERENCES

- Cooray, V., M. Rahman, V. Rakov, On the NOx production by laboratory electrical discharges and lightning, *J. of Atmo. Solar-Terres. Phys.*, **71**, 1877-1889, 2009.
- Koshak, W. J., R. J. Solakiewicz, R. J. Blakeslee, S. J. Goodman, H. J. Christian, J. M. Hall, J. C. Bailey, E. P. Krider, M. G. Bateman, D. J. Boccippio, D. M. Mach, E. W. McCaul, M. F. Stewart, D. E. Buechler, W. A. Petersen, D. J. Cecil, North Alabama Lightning Mapping Array (LMA): VHF source retrieval algorithm and error analyses, *J. Atmos. Oceanic Technol.*, **21**, 543-558, 2004.
- Koshak, W. J., M. N. Khan, A. P. Biazar, M. Newchurch, R. T. McNider, A NASA model for improving the lightning NOx emission inventory for CMAQ, Joint Session: 4th Conference on the Meteorological Applications of Lightning Data and the 11th Conference on Atmospheric Chemistry; 89th Annual American Meteorological Society (AMS) Conference, Phoenix, AZ, January 11-15, 2009.
- Koshak, W. J., H. S. Peterson, E. W. McCaul, A. Biazar, Estimates of the lightning NO_x profile in the vicinity of the North Alabama Lightning Mapping Array, International Lightning Detection Conference (ILDC), Orlando, FL, April 19-20, 2010.
- Koshak, W., H. Peterson, M. Khan, A. Biazar, L. Wang, The NASA Lightning Nitrogen Oxides Model (LNOM): Application to Air Quality Modeling, XIV International Conference on Atmospheric Electricity (ICAE), Rio de Janeiro, Brazil, August 8-12, 2011.
- Koshak, W. J., and H. S. Peterson, A summary of the NASA Lightning Nitrogen Oxides Model (LNOM) and recent results, 10th Annual Community Modeling and Analysis System (CMAS) Conference, Chapel Hill, NC, October 24-26, 2011.
- Koshak, W. J., H. S. Peterson, A. P. Biazar, M. Khan, and L. Wang, 2013: The NASA Lightning Oxides Model (LNOM): application to air quality modeling, *Atmos. Res.*, <http://dx.doi.org/10.1016/j.atmosres.2012.12.015>.
- Wang, Y., A. W. DeSilva, and G. C. Goldenbaum, 1998: Nitric oxide production by simulated lightning: dependence on current, energy, and pressure, *J. Geophys. Res.*, **103**, 19149-19159.