











# AAS WorldWide Telescope: A Seamless, Cross-platform Data Visualization Engine for Astronomy Research, Education, and Democratizing Data

Philip Rosenfield<sup>1</sup> , Jonathan Fay<sup>1</sup> , Ronald K Gilchrist<sup>1</sup> , Chenzhou Cui<sup>2</sup> , A. David Weigel<sup>3</sup> , Thomas Robitaille<sup>4</sup> ,  
Oderah Justin Otor<sup>1</sup> , and Alyssa Goodman<sup>5</sup> 

<sup>1</sup>American Astronomical Society, 1667 K St NW Suite 800, Washington, DC 20006, USA; [philip.rosenfield@aaas.org](mailto:philip.rosenfield@aaas.org)

<sup>2</sup>National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing, 100012, People's Republic of China

<sup>3</sup>Christenberry Planetarium, Samford University, 800 Lakeshore Drive, Birmingham, AL 35229, USA

<sup>4</sup>Aperio Software Ltd., Headingley Enterprise and Arts Centre, Bennett Road, Leeds, LS6 3HN, UK

<sup>5</sup>Harvard Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

Received 2017 October 25; revised 2018 January 24; accepted 2018 January 24; published 2018 May 11

## Abstract

The American Astronomical Society's WorldWide Telescope (WWT) project enables terabytes of astronomical images, data, and stories to be viewed and shared among researchers, exhibited in science museums, projected into full-dome immersive planetariums and virtual reality headsets, and taught in classrooms, from middle school to college. We review the WWT ecosystem, how WWT has been used in the astronomical community, and comment on future directions.

*Key words:* astronomical databases: miscellaneous – atlases – catalogs – surveys – virtual observatory tools

## 1. Introduction

WorldWide Telescope<sup>6</sup> (WWT) was first designed as a tool for astrophysical data exploration and discovery in the era of large and disparate data (Gray & Szalay 2004). WWT was an early attempt to answer an emerging question: “how will research techniques change in the era of multi-wavelength, multi-epoch, large data sets hosted throughout the web?” Launched in 2008 by Microsoft Research, WWT was open source (under the MIT license) seven years later with copyrights transferred to the .NET Foundation and project direction and management taken up by the American Astronomical Society (AAS). Recent and near future AAS WWT development is focused on implementing operating-system-agnostic features to better enable astronomers to use WWT either individually or through archival data centers.

WWT is a scientific data visualization platform (see Figure 1). It acts as a virtual sky, allowing users to explore all-sky surveys across the electromagnetic spectrum, to overlay data from NASA's Great Observatories, and to import their own imagery and tabular data. In WWT's 3D environment, users can explore planetary surfaces and elevation maps, orbital paths and ephemerides of major and minor solar system bodies and asteroids, and zoom-out from the solar system to view the positions and approximate colors of stars from the HIPPARCOS catalog (Perryman et al. 1997) and galaxies of the Cosmic Evolution Survey data set (COSMOS; Scoville et al. 2007).

WWT's unique contextual narrative layer (Wong 2008) has set it apart from the myriad other available data visualization systems. Users can record or view recorded paths, though the virtual environments add narration, text, and imagery. With these “tours” users can share their stories, outreach specialists can create programming, and researchers can distill their discoveries to other researchers or to the public.

These features make WWT a seamless data visualization program with an engaging learning environment. This combination has enabled WWT to deliver astrophysical data hosted around

the world not only to researchers, but to classrooms, planetariums, and educational outreach programs. Therefore, WWT is more than a tool to aid astronomical research and discovery; it helps to democratize astronomical data and knowledge by enabling equal data access for all users. For example, only a few dozen content experts across academia and science centers have recorded tours curated within WWT or have organized WWT-related outreach efforts, yet WWT has reached millions<sup>7</sup>: from explorers using the WWT application; to students visiting planetariums and museums around the world; to students at all levels using WWT to augment their curricula. WWT may be the only visualization software that when astronomers add data and context, it becomes available to students—who may be geographically distant from a science center—to also investigate.

We overview the technical aspects of WWT in Section 2, describe some major uses of this visualization tool in Section 3, outline the open source management and sustainability plan in Section 4, and conclude in Section 5.

## 2. The WWT Ecosystem Overview

In the nearly 10 years of WWT development, several applications, services, and technologies have been created in order to deliver astronomical data from around the globe in a seamless visualization and narrative environment. In this section, we outline the main components of the WWT software ecosystem.


### 2.1. Clients

WWT is offered as a standalone Microsoft Windows application and web-based applications and services. All of the code base is open source under the MIT license and is hosted on GitHub.<sup>8</sup>

<sup>7</sup> Over 10 million unique Windows Application downloads from 2007 to 2014.

<sup>8</sup> See the WWT meta-repository: <https://github.com/WorldWideTelescope/wwt-home>.

<sup>6</sup> <http://worldwidetelescope.org>



**Figure 1.** Screen captures of the WorldWide Telescope User Interface. Clockwise from top left: “Sky” mode showing *IRIS: Improved Reprocessing of Infrared Astronomy Satellite (IRAS) Survey* (12, 60, and 100  $\mu\text{m}$  (Red is IRIS100, green is IRIS60, blue is IRIS12) cross-faded against the 100  $\mu\text{m}$  David J. Schlegel, Douglas P. Finkbeiner and Marc Davis (SFD) Dust Map (data are provided by *IRAS* and the *Cosmic Background Explorer*, Princeton University, and University of California, Berkeley. Original *IRAS* data: NASA/JPL IPAC, *IRIS* Reprocessing: Canadian Institute for Theoretical Astrophysics/Institut d’Astrophysique Spatiale. TOAST-formatted mosaics were obtained using the facilities of NASA’s SkyView Virtual Telescope.); “Planet” mode showing Mars visible imagery (NASA/USGS/Malin Space Science Systems/JPL); “3D solar system” mode (which includes HIPPARCHOS and COSMOS data) showing planets and their orbital paths, Oort cloud objects (white), Jupiter Trojans (yellow), and main belt asteroids (purple to white, colored by zone). Minor Planet ephemerides from the International Astronomical Union’s Minor Planet Center); and “Panorama” mode showing an image taken by Apollo 12 Lunar Module Pilot Alan Bean in 1969 and assembled at the NASA Ames Research Center in 2007. (Images are provided courtesy of NASA and the Lunar and Planetary Institute).

### 2.1.1. Windows Application

The Microsoft Windows application contains the richest set of features and customizations. It currently serves as the basis of features that are ported to Web application. The WWT Windows application can run natively or through virtualization on either a local virtual machine or on the Cloud.

Beyond the main tasks of visualizing the sky and 3D universe, the WWT Windows application features flexible display modes, supports a variety of controllers, and can incorporate a wide variety of data.

WWT can be displayed in power walls (ultra-high-resolution displays created by connecting several monitors or projectors), 3D-stereo projections, virtual reality headsets, interactive kiosk exhibits, and multi-channel environments (e.g., in multiple-projector full-dome planetariums). Due to the high demand on computational resources, there are limited plans to port these flexible display modes to the web application in the near future.

Users may drive WWT with their keyboard and mouse or customize a MIDI controller, an XBOX controller, or the Microsoft Kinect.

A wide variety of data layers are both customizable and controllable. Users can import 3D Models, tabular data, orbital data, imagery (including all-sky maps and planet surface maps), planetary digital elevation models, and catalogs. We discuss the WWT data files in more detail in Section 2.5.

### 2.1.2. Web Application

A growing subset of features from the WWT Windows application are now delivered via a plugin, free, browser-based platform rendered by WebGL. The web application uses the same curated image and catalog data as the Windows application (see Section 2.5), it supports built-in tour authoring and playback, including a simple slide-based or complex key-framed animations. The web application can display tiled images, tours with audio and imagery, and most of the WWT intrinsic data layers.

The WWT Web application leverages the Web Control API for rendering. To better serve astronomical researchers, current development efforts led by the American Astronomical Society are focused on porting more data layer features of the Windows application to the web application and exposing them in the Web Control API (see Section 2.3).

### 2.1.3. Extensibility of WWT Clients

Users can extend and customize their WWT client experience at a variety of experience levels. They can create a slide-based tour with linked slides (i.e., similar to a DVD menu). With the Windows application users can have full control over WWT tour rendering using the time-line function (see [this guide](#)). They can add their own data layers (see Table 1 for a list of supported formats). And they can configure

**Table 1**  
Supported Data Formats

Name	Description	Reference or Documentation
WTML (also called a “Collection File”)	XML-coded metadata description of how to find places and streaming data services.	<a href="#">Collections Guide</a>
Tiled Data	Tiled (quad-tree) multi-resolution images sets in a variety of projections delivered through HTTP streams and described by metadata in a WTML file. <i>Supported Projections</i> 1. TOAST: Tessellated Octahedral Adaptive Subdivision Transform (TOAST; T. McGlynn et al. 2018, in preparation) is a Hierarchical Triangular Mesh (HTM; Kunszt et al. 2000) based spherical projection without singularities at the poles and delivered through a quad-tree tile structure. 2. Mercator: Mercator-projected tiles (such as those provided by Bing and Google maps), for all-sky data; the poles are cut off. 3. Equirectangular (plate carrée): equidistant cylindrical projection for all-sky images; the poles have a singularity. 4. Tangential Study Maps: TAN-projected sections of small areas of the sky.	<a href="#">Projection Reference</a>
VAMP	VAMP-encoded images can be loaded and viewed directly, or tiled by our web service.	<a href="#">Virtual Astronomy Multimedia Project Standards Guide for Adding 3D Models</a>
3D models	.3ds or .obj 3D file formats including textures and lighting.	...
Tabular Data	Either through cut/paste or import, tabular data can be visualized through an interactive user interface on the Windows application.	
WMS	Web Mapping Service source for tiled and time-series data.	<a href="#">Guide for adding WMS Data</a>
VO Table	Tabular Data in VO Table Format can be viewed as tables or visualized	<a href="#">VOTable Format Definition</a>
FITS	Flexible Image Transport System (FITS) files in TAN-projection with the World Coordinate System (WCS) in celestial coordinates can be viewed and stretched.	<a href="#">FITS and WCS Standard</a>
Shapefiles and WKT	Shape files can be loaded and viewed and converted to Well Known Text (WKT) and Tables. WKT can be included in tabled for viewing complex geometry along with a text extension for displaying oriented text in 3D.	<a href="#">ESRI Shapefile Technical Description</a>
ODATA	Tabular feeds from Open Data (ODATA) sources can be mapped in layers and dynamically refreshed at load time.	<a href="#">Open Data Specification</a>
TLE	Two Line Element orbital data can be used for bulk display of orbits, or as a foundation for a reference frame.	<a href="#">Guide for adding TLE Data Definition of TLE</a>
SPICE (Acton 1996; Acton et al. 2017)	NASA SPICE kernel data can be imported to display spacecraft or planetary ephemerides by minor manipulation within the NASA SPICE GUI and Microsoft Excel into an .xyz text file.	<a href="#">Guide for adding NASA SPICE Data About SPICE</a>

a planetarium with multiple computers and projectors to display WWT (see [this guide](#)). Below we mention two more means to extend the WWT clients, by using “communities” and by contributing documentation.

*Communities*—user-uploaded WWT content (e.g., tours and data layers) that is available to all users of the WWT clients through the Communities tab. Communities content can also be viewed on the WWT website without launching the web application (see [this guide](#)).

*Documentation*—all WWT documentation is now user-contributed and maintained on GitHub.<sup>9</sup> Each guide is rendered from Markdown as a GitBook,<sup>10</sup> which is available to read online or downloadable in PDF and eBook formats. There are also a growing number of user-contributed WWT tutorials on YouTube. WWT and the WWT Ambassadors (see Section 3.1) maintain YouTube channels ([WWT Videos](#), [WWT Ambassadors Videos](#)). We further discuss contributing documentation in Section 4.

## 2.2. Web Services

*Embeddable Web Control*<sup>11</sup>—crosses the line between a flexible web application and a software development kit

<sup>9</sup> See the WWT documentation meta-repository: <https://github.com/WorldWideTelescope/wwt-documentation>.

<sup>10</sup> <http://gitbook.com/>

<sup>11</sup> Available at <http://worldwidetelescope.org/Use/Embed>.

(SDK). It implements a (growing) subset of the WWT Web user interface to support visualizing data sets or playing tours. It is designed to be cut-and-pasted into websites outside of the [worldwidetelescope.org](http://worldwidetelescope.org) domain.

The embeddable code-generation tool creates a WWT namespace on the containing page and spawns an `iframe` that hosts the control. The `webcontrol.aspx` script reads from the options in the containing `div` and passes them to the `iframe` via a `postMessage` API (for cross-domain scripting).

*Show Image Service*—enables an image to be shared by embedding information about the image in a URL; see the [documentation](#) for more details and Table 2 for examples of its usage.

## 2.3. APIs

WWT has several application programming interfaces (APIs) of interest to Web and Windows developers. While all of the WWT code base is open source, it is incredibly complex, and above the software engineering level of most astronomers and astronomical software engineers. We expect most developers interested in extending or customizing WWT to do so through the APIs (see Section 4 for more discussion on code contributions).



**Table 2**  
Show Image Service Usage Examples

Name, Website	Instructions
<a href="#">Montage</a>	Instructions on using Montage to TOAST images for WWT.
<a href="#">Chandra Photo Album</a>	Click on an image (from 2012 or earlier), then click on the right panel “View on the Sky (WWT).”
<a href="#">Hubble Space Telescope</a>	Click on an image and scroll to the bottom right “View in WorldWide Telescope.”
<a href="#">Astrometry.net</a>	Click on or solve an image and scroll to the bottom right “View in WorldWide Telescope.” (Also available as a Flickr plugin.)
<a href="#">Spitzer IPAC AstroPix</a>	Click on an image and click “View in WorldWide Telescope” on the top right.

**Table 3**  
Web Control API Examples

Name, Website	Description	Supplementary Code
<a href="#">Import Image</a>	Import an AVM (Astronomy Visualization Metadata Standard) tagged image or solve for AVM tags with <a href="#">astrometry.net</a> and display it in the WWT Web Application.	<a href="#">ImportImage.js</a>
<a href="#">Great Observatories</a>	NASA Great Observatories overview and image interactive.	<a href="#">Observatories.js</a>
<a href="#">Planet Explorer</a>	Travel from planet to planet in the 3D solar system mode.	<a href="#">PlanetExplorer.js</a>
<a href="#">Spectroscopy</a>	Spectroscopy overview and interactive.	<a href="#">Spectroscopy.js</a>
Example Web Control API Usage Around the Web		
<a href="#">GLIMPSE 360 WWT Viewer</a>	“An infrared view of the disk, or plane, of the Milky Way galaxy assembled from more than 2 million snapshots taken over the past 10 years by NASA’s <i>Spitzer Space Telescope</i> .”	...
<a href="#">Interactive Planck Viewer</a>	“...Compare the various components of the sky derived from <i>Planck</i> observations, as well as some other all-sky data sets.”	...
<a href="#">ADS All Sky Survey</a>	“The ADS All-Sky Survey is an ongoing effort aimed at turning the NASA Astrophysics Data System (ADS), widely known for its unrivaled value as a literature resource for astronomers, into a data resource.”	<a href="#">ADSASS WWT-frontend</a>
<a href="#">Milky Way 3D</a>	“A tool intended to organize and curate links to information about data sets relevant to our 3D understanding of the Milky Way.”	...
<a href="#">Chandra Source Catalog Release 2.0</a>	The <i>Chandra</i> Source Catalog (CSC) is ultimately intended to be the definitive catalog of X-ray sources detected by the <i>Chandra X-ray Observatory</i> . To achieve that goal, the catalog will be released to the user community in a series of increments with increasing capability. The first official release of the CSC 2.0 will include 10,382 <i>Chandra</i> ACIS and HRC-I imaging observations released publicly through the end of 2014. When complete, CSC 2.0 will provide information for about 370,000 unique sources in several energy bands.”	...
<a href="#">pyWWT</a>	The pyWWT package aims to make it easy to use WorldWide Telescope from Python (see Section 3.4).	<a href="#">pyWWT on GitHub</a>

### 2.3.1. Windows APIs

The *Layer Control API (LCAPI)*—allows local or remote applications to send data and control data layers in the Windows application for integration in scenarios outside of the [worldwidetelescope.org](#) domain. There is deep integration with the LCAPI and Microsoft Excel through an Excel Add-In tied to the LCAPI, as well as python bindings (see Section 3.4).

*Windows Client Socket API*—is open source, and used by WWT Kinect control to interact with WWT.

### 2.3.2. Web APIs

The *Web Control API*—is a complex and robust engine that renders WWT objects and imagery into a canvas element. Its purpose is to deeply integrate WWT visualization and tour playback into websites outside of [worldwidetelescope.org](#). It was written in C# and compiled into javascript using scriptsharp (See the [API Documentation](#)). Table 3 lists examples of Web Control API usage.

*Web Services API*—sets of hosted web services that deliver or process data to and for the WWT clients and other

consumers, such as the ShowImage service referenced in Table 2. Related documentation is hosted on [GitHub](#).

### 2.4. Tools and SDKs

While part of Microsoft Research, WWT published SDKs to assist users and developers with importing data (available [here](#)). These tools are in the process of being converted to web services to support all operating systems.

*Sphere Toaster*—takes small to medium sized images and outputs Tessellated Octahedral Adaptive Subdivision Transform (TOAST; T. McGlynn et al. 2018, in preparation) tile pyramids, plate files, and WTML.

*Study Chopper*—takes small to medium study images and outputs TAN-projected tile pyramids and WTML.

*AVM Import Tool*—takes Astronomy Visualization Metadata Standard-tagged images with WCS coordinates and creates hosted tile pyramids and WTML files to access them. This service is available online [here](#).

*Tile SDK*—A software tool kit with samples that allows user to create a custom data transformation pipeline to take images or elevation data from input format and output it to WWT-compatible pyramids.

*Terapixel Project*—while not exactly an SDK, the Terapixel Project is an open source pipeline that was used to reprocess the original DSS data from scanned plates into a Terapixel TOAST pyramid with globally optimized color and brightness corrections (Agarwal et al. 2011).

### 2.5. Curated Data

The data shown in WWT are a mix of curated data (stored on the Azure Cloud) and data displayed from around the globe, including the National Virtual Observatory (NVO) registries. There are currently more than 90 all-sky surveys, 20 planet maps, and 60 panoramas (see Tables 4–6 in the Appendix). In addition, WWT hosts over 500 contributed images from education and public outreach offices at Hubble, *Chandra*, *Spitzer*, Gemini, NOAO, and NRAO, as well as from astrophotographers.

### 2.6. Supported Data Formats

WWT supports a variety of data formats (see Figure 2), which users can import via the Windows application. Importing of data on the web application is in development, and natively it currently supports FITS images, and many more data formats are possible through an astropy-affiliated package (see Section 3.4). Table 1 lists current levels of data support.

#### 2.6.1. Plate Files

The reasoning behind splitting large images into tiles is to read from disk-only as much as is absolutely necessary for each database query (e.g., a view of a portion of the sky). The efficiency that tiling enables gives way in the limit of reading and transferring many small files from the server. The WWT team created “plate files” as an efficient way to store, exchange, and deliver tiled multi-resolution images. Plate files have an index and data storage and allow millions of small files to be packed into a single file. Single large files copy up to two orders of magnitude faster than the millions of files they contain. There exist two plate file formats:

1. For densely packed complete surveys or studies with single-generation data.
2. For sparse data that can contain multiple generations of data for a given tile index, this format is great for building large surveys over time with streaming data, such as, for example, the Mars HiRise data.

## 3. WWT Usage

WWT has been used extensively at nearly all levels of astronomy education. For clarity, we separate the following section by formal and informal education, and include education and public outreach (EPO) centers as part of informal education.

### 3.1. Formal Education

The most prominent formal education program that uses WWT is the WWT Ambassador Program<sup>12</sup> (WWTa), which has evolved alongside WWT since 2010.

The WWT Ambassadors Program aims to educate the public about astronomy and science using WWT. It is run by a team of

astronomers and educators at Harvard University, in collaboration with the AAS and Microsoft Research. They recruit and train volunteer ambassadors, who help facilitate the use of WWT in educational settings like schools, science festivals, and museums.

#### 3.1.1. Pre-college Courses

WWTa have directly served thousands of students in the classroom. Since 2015, more than 700 of these students had extensive learning experiences that included one or two full weeks of instruction from WWTa team members (for more about Moon phases, seasons, and life in the universe, see Udomprasert et al. 2014, 2016). They have pre-post assessment data showing outstanding learning gains from these interventions (Udomprasert et al. 2012, P. Udomprasert et al. 2018, in preparation). Lesson plans and other materials are available on their website.<sup>13</sup>

WWT has also been used for Astronomy classes at Beijing Shijia Primary School for three years. A guidebook on Interactive Astronomy Teaching (Primary School Teacher version), designed by the Chinese Virtual Observatory project, was published by Popular Science Press in Beijing.

#### 3.1.2. College Courses and Beyond

*Astronomy Courses*—the WWTa program has helped to create WWT Windows application-based introductory astronomy labs on parallax and Hubble’s law. They have been used at Bucknell University and the University of Massachusetts Amherst (see Ladd et al. 2015, and the [WWTa Website](#)).

WWT has been part of introductory astronomy courses at the Central China Normal University since 2009 (Qiao et al. 2011; Wang et al. 2015), and a graduate ISM course at Harvard University (Sanders et al. 2014). Co-author Weigel uses WWT to visualize spacecraft ephemerides and planetary texture maps in context in an immersive full-dome environment for the Introduction to Astronomy class at Samford University. These visuals are presented and manipulated in real time in the Christenberry Planetarium on Samford’s campus.


*General Science Courses*—Samford University makes further use of WWT in their Cultural Perspectives and Scientific Inquiry classes. Weigel teaches about the Scientific Revolution using WWT in a planetarium. Students are guided on a journey from naked-eye astronomical observations and antiquated methods of scientific reasoning and philosophy, through the age of the scientific revolution and the subsequent invention of telescopes, and finish with a look at current big data astronomy and predictions for the future.

*Science Communication*—The University of Washington offered an undergraduate seminar on science communication as applied to creating a tour in WWT (the curriculum is available here<sup>14</sup>). Students learned storyboarding, distilling versus dumbing down, avoiding jargon, and how to create WWT tours playable in the University of Washington planetarium.

<sup>13</sup> <https://wwtambassadors.org/science-education-research>

<sup>14</sup> [http://philrosenfield.github.io/teaching\\_files/tourmaking/CourseGuide.pdf](http://philrosenfield.github.io/teaching_files/tourmaking/CourseGuide.pdf)

<sup>12</sup> <https://wwtambassadors.org>



**Figure 2.** Examples of supported data and models. Clockwise from top left: view of Earth with near-Earth objects imported as Two Line Element (TLE) files (data from International Astronomical Union’s Minor Planet Center); 3D ISS Model (ISS Model from: Toshiyuki Takahe); a 3D rendering of a user-plotted catalog (the CfA Redshift Survey, green; Geller & Huchra 1989) over the COSMOS data set (Scoville et al. 2007); latitude, longitude, and altitude of a Mars Rover imported to WWT using the WWT-Excel plugin (data courtesy of Joe Knapp, [curiosityrover.com](http://curiosityrover.com)).

### 3.2. Informal Education

WWT has been shown to be an excellent data-driven astronomical education and public outreach environment. For example, it inspired co-author Cui to initiate the IAU Working Group “Data driven Astronomical Education and Public Outreach” (DAEPO; Cui & Li 2018), it has been used to help capture and share indigenous Australian astronomical knowledge (Nakata et al. 2014), and has allowed planetariums to be “flipped,” allowing students to present planetarium shows to their peers.

#### 3.2.1. Museums and Planetariums

WWT has been used for live and recorded shows in major planetariums, such as the Adler Planetarium (Cosmic Wonder;<sup>15</sup> Figure 3) and the Morrison Planetarium (Earthquake).<sup>16</sup>

WWT has also proven to be an attractive choice for smaller planetariums, since the software is free and it comes with a robust projector calibration system.<sup>17</sup> The following list of WWT-driven smaller planetariums is incomplete, but includes the University of Washington, Mount Hood Community

College, Central Washington University, Samford University, and Bellevue College. In China, six WWT-driven planetariums have been built and two are currently under construction, with an Internet-based WWT-driven planetarium alliance for resource and experience exchange.

Mobile, portable, or traveling planetarium programs have also begun to use WWT. For example, Discovery Dome<sup>18</sup> packages WWT software with their inflatable domes, and the University of Washington, the University of Oklahoma (as the Soonertarium), and Harvard University all have WWT-driven mobile planetariums programs. Nearly every mobile planetarium program has participated in some form of a local science festival; for example, Figure 4 shows students watching a tour in a portable WWT planetarium.

One of the unique uses of WWT in an educational setting comes from “flipping” the classroom or in this case, the planetarium. Instead of students passively attending a lecture or recorded video in a planetarium, they can research and then create their own content to share in a planetarium. This application of WWT has been independently described by several groups. The DAEPO has coordinated student-authored WWT tour contests<sup>19</sup> three times. The Cal Academy helped

<sup>15</sup> <http://www.adlerplanetarium.org/events/cosmic-wonder/>

<sup>16</sup> <https://www.calacademy.org/press/releases/earthquake-planetarium-show-opens-on-may-26-2012-at-the-california-academy-of>

<sup>17</sup> See the Multi-Channel Dome Guide at <https://legacy.gitbook.com/book/worldwidetelescope/worldwide-telescope-multi-channel-dome-setup/details>

<sup>18</sup> [http://eplanetarium.com/software\\_worldwide\\_telescope.php](http://eplanetarium.com/software_worldwide_telescope.php)

<sup>19</sup> [http://english.nao.cas.cn/ns/ConferenceNWorkshop/201706/t20170614\\_178121.html](http://english.nao.cas.cn/ns/ConferenceNWorkshop/201706/t20170614_178121.html)





**Figure 3.** Adler Planetarium Audience viewing WWT during Cosmic Wonder.



**Figure 5.** WorldWide Telescope rendered in the full-dome Christenberry Planetarium at Samford University.



**Figure 4.** 2016 Science and Technology Week (China). Children are watching a WWT tour shown inside a portable planetarium.



**Figure 6.** Samford University student experiencing VR over the surface of Pluto (New Horizons) in WWT.

San Francisco Bay Area students to present WWT in the Morrison Planetarium (Roberts 2014). The University of Washington works with Seattle-area middle and high schools to present WWT in a mobile planetarium (Rosenfield et al. 2014). The Christenberry Planetarium at Samford University mentors middle and high school students in summer programs, teaching them to produce and present WWT tours for public audiences in a planetarium (see Figure 5).

The Christenberry Planetarium also uses WWT’s VR capabilities in public outreach to promote the latest astronomical discoveries and research by adding the “wow” factor using cutting-edge technology within the university and the greater Birmingham, Alabama community. Weigel and his students develop VR specific WWT tours and host programs to both inspire and teach youth in astronomy in Alabama (see Figure 6).

WWT has a kiosk mode where a user can interact with a guided tour or contained aspect of WWT. An incomplete list of museums and informational centers that have used WWT in this way include Harvard University (see Figure 7), the Adler Planetarium (see the Space Visualization Lab<sup>20</sup>), and the Imiloa

Astronomy Center of Hawai’i, which had a Microsoft Kinect driven WWT exhibit “The Universe at your Finger tips”.<sup>21</sup>

### 3.2.2. Education and Outreach Centers

Both WWT clients have served as a conduit for publishing EPO imagery and stories. For example, the “Explore” tab shows curated feeds from the *Hubble Space Telescope*, *Spitzer* Science Center, *Chandra X-ray Observatory*, and the European Space Agency. Several outreach efforts have also made use of the WWT Web Control API (see bottom of Table 3) and the WWT Show Image service (see Table 2).

### 3.3. Astronomical Research

WWT enters the astronomical research workflow at several points: (1) discovery, helping visualize large and wide data sets; (2) analysis, WWT APIs allow analysis tools to link data; and (3) dissemination, WWT can enhance figures and presentations, and researchers can create video abstracts to accompany their submitted manuscripts to journals (see Section 3.5). WWT and its APIs allow observatories and mission centers to adopt a common visualization interface for

<sup>20</sup> <http://www.adlerplanetarium.org/whats-here/dont-miss/space-visualization-lab/>

<sup>21</sup> [https://subarutelescope.org/Gallery/movie\\_subaru.html](https://subarutelescope.org/Gallery/movie_subaru.html)




Figure 7. Harvard University's WWT-driven Kiosk.

their databases. In the following section, we discuss the most recent efforts and future outlook led by the AAS to help the astronomical research community benefit from WWT technologies.

### 3.4. Discovery and Analysis

Astronomers interested in using survey data (e.g., SDSS, LSST, *Gaia*, DESS) have most likely shifted from the traditional *ftp-grep* data analysis model, where a scientist first downloads a copy of data (*ftp*) and then uses an analysis package to identify for patterns, or answer their research question (*grep*) (see, e.g., Gray & Szalay 2004). Instead, astronomers analyze their data by querying remote databases, and soon, accessing hosted computing facilities to perform their analysis (e.g., LSST: Jurić et al. 2015, and refs. therein). WWT was first designed to exist in the post *ftp-grep* regime, and now that it has been open source and ported to the web, it has plug-and-play potential for use in smaller-budgeted observatories and data archives.

*Archival Centers*—the next release of the WWT Web Control API will have data import/export, and selection features that will make it a powerful link between data archives and researchers. Observatories and data centers can insert the web control on their website and have a robust, open source, data archive visualization engine. Observatories or data archive centers that use WWT in this way will automatically benefit from their EPO efforts, since EPO staff would have the same access to data and be able to create tours or build a custom user interface over the WWT Web Control API.

In addition, abstract services such as ADS<sup>22</sup> can use the WWT interface as another means for users to discover research in astronomy. The Zooniverse's Astronomy Rewind project<sup>23</sup> is a citizen scientist project to place AAS journal figures dating back to the 1800s into the Astronomy Image Explorer<sup>24</sup> with AVM tags so they can be viewed in WWT.

*Data Analysis Tools*—several data analysis tools are being built off of the Web Control API. For example, a link between JS9<sup>25</sup> and WWT has been established as part of the AAS

<sup>22</sup> <http://adsabs.harvard.edu>

<sup>23</sup> <https://www.zooniverse.org/projects/zooniverse/astronomy-rewind>

<sup>24</sup> <http://www.astroexplorer.org>

<sup>25</sup> <https://js9.si.edu>: the online javascript version of DS9 (<http://ds9.si.edu>) with a public API.




Figure 8. A screenshot of pyWWT, a Python interface for the WorldWide Telescope based on the Web Control API.

Astrolabe project.<sup>26</sup> Users of this service can upload their images to JS9, manipulate them as they would in DS9, and view the result in WWT.

Python bindings for the WWT Web Control API are a new focus and were first prototyped in Glue-viz.<sup>27</sup>

*PyWWT: A Python interface to WWT*—PyWWT is a package that provides access to much of the Windows and web clients' functionality to Python on a variety of platforms. It allows the WWT web client to be used as an interactive widget inside Jupyter Notebooks and Jupyter Lab (pictured in Figure 8; Kluver et al. 2016), as well as through a standalone interactive Qt-based viewer/widget,<sup>28</sup> and also provides a Python-based client for the WWT desktop application on Windows (using the LCAPI, see Section 2.3).

In all cases, users can pan and zoom with the mouse, display multiple layers from all-sky surveys, and play tours. PyWWT makes it easy to add annotations, or shapes, that can be used to highlight celestial objects, enhance tours, or illustrate fields of view for telescopes, among many other applications. The package is written to integrate smoothly with Astropy (Astropy Collaboration et al. 2013) for easy adaptability to typical research workflows such as those described in Section 3.4. Support for Astropy tables and displaying local FITS images is forthcoming.

PyWWT is under active development and is hosted under the WorldWideTelescope GitHub organization (see Table 3), where its full code base is open source (version 0.3.0 was launched in 2017 December).

### 3.5. Dissemination

We have seen examples of the WWT narrative layer (tour features) in sharing stories and teaching astronomy. WWT has also been used in video abstracts, which have converted WWT tours into videos distilling a scholarly publication (see the Publisher's website of Currie et al. 2015; Yusef-Zadeh et al. 2015;

<sup>26</sup> The Astrolabe Project ([astrolabe.arizona.edu/](http://astrolabe.arizona.edu/)) is creating a new open-access repository for previously un-curated astronomical data sets, building on existing CyVerse infrastructure with robust cloud-based resources for managing, linking, processing and sharing research data.

<sup>27</sup> Glue-viz is a Python library for exploring relationships within and among related data; see <http://glueviz.org/en/stable/>, and see the WWT Plugin at <http://glueviz.org/en/stable/whatsnew/0.11.html?#experimental-worldwide-telescope-plugin>.

<sup>28</sup> <https://www.qt.io>



Batygin & Brown 2016; Berriman & Good 2017). Future WWT Web Control development will allow for embedding WWT in 2D and 3D modes as interactive figures in scholarly publications.

#### 4. Management and Sustainability of WWT

WWT is managed by the AAS and its (MIT) copyright is held by the .NET Foundation. Below we overview the management and sustainability plan for this open source software project.

##### 4.1. Project Leadership

AAS Board of Trustees (AASBoT) is in charge of managing the AAS WWT project. The AASBoT has mandated that the AAS WWT Steering Committee continually assess the WWT project against metrics based on the WWT's purpose statement, as well as the AAS' purpose statement. The AAS WWT Steering Committee and AAS WWT Project Director work together to interpret the purpose statements into initiatives with quantitative metrics for success.

##### 4.2. Open Source Contributions

The longevity and success of open source software depend on the community. We are striving to build an inclusive WWT community. We expect everyone in the WWT community to follow our code of conduct and a contributing guide. Below, we discuss two avenues of contribution: contributing to the code base, and contributing documentation and examples.

###### 4.2.1. Code Contributions

We maintain a meta-repository, `wwt-home` ([github.com/WorldWideTelescope/wwt-home](https://github.com/WorldWideTelescope/wwt-home)), which primarily contains a table of information on the different WWT-related repositories in the WorldWideTelescope GitHub organization. Importantly, each table entry includes a description of the project as well as the skills required to contribute. Until we have a robust user community to review code contributions (as well as proposals for enhancements), code contributions will be considered by the WWT Advisory Board.

###### 4.2.2. Documentation Contributions

As discussed in Section 2.1.3, all of our documentation has been open source and hosted on GitHub, accessible via our meta-repository, `wwt-documentation` ([worldwidetelescope.github.io/wwt-documentation/](https://worldwidetelescope.github.io/wwt-documentation/)) and including code samples.

Contributors can edit existing documentation or create their own, even if they are not familiar with GitHub or Markdown. In addition, they can create documentation within a GitBook or import files created using Microsoft Word. Each new contribution will be added as a line to the `wwt-documentation` table, which is the basis of the WWT website's User Guides Page.

Sample code can be published as gists and linked to our meta-repositories or as Jupyter notebooks (e.g., `pyWWT` example notebooks).<sup>29</sup>

We will continue to engage and consult with the broader astronomical community to plan new features and services via AAS member polls, the WWT online forum, tracking GitHub

issues, the WWT help desk ([wwt@aaas.org](mailto:wwt@aaas.org)), and feedback during workshops and webinars (direct and via assessments).

##### 4.2.3. Contributing WWT Products

In the future, we hope to create a place on the web to house WWT-related workshop materials, contributed WWT lesson plans, 3D models, music, and more—that meet the existing skill sets of the contributors.

#### 5. Conclusions

We have reviewed the software ecosystem of WWT and many of its uses in astronomy. WWT is astronomical visualization software that can directly link to all levels of education. In that way, WWT could democratize astronomical data, but only with the help of the astronomical community.

In summary:

1. WWT is open source (MIT license) and managed by the American Astronomical Society.
2. WWT has two main applications: (1) a Microsoft Windows application that is a mainstay in planetariums and international education; and (2) a WebGL-powered Web application that is operating-system-agnostic, which we expect most US-based astronomers to prefer.
3. Each application has APIs and services to allow users and developers to customize and extend WWT without needing the high level of C# programming knowledge to advance the core WWT components.
4. WWT has been used innovatively in formal and informal educational settings, supplementing: (a) early learners' physical understanding of the seasons and moon phases; (b) college students' introductory astronomy and general science courses, as well as at least one graduate astronomy course; (c) planetarium education, e.g., by enabling students to present planetarium shows to their peers.
5. Educational programs have used WWT in a variety of settings: computer labs, kiosks, virtual reality headsets, and planetariums.
6. Ongoing development of the WWT Web Control API will help to embed the powerful tool in astronomical research: (a) as a visualization engine for data archives; (b) as a link to python and javascript analysis tools; and (c) as interactive (or video) figures for scholarly publications and presentations.

The authors thank the WorldWide Telescope Advisory Board, whose efforts to support the WWT project since its inception and help to transition WWT from Microsoft to the AAS, ensured its existence. We specifically thank Sarah Block, Andy Connelly, [Chen-Zhou Cui](#), Cristine Donnelly, Karl Fogel, [Ron Gilchrist](#), [Alyssa Goodman](#), Morgan Griffith, [Bryan Heidorn](#), [Robert Hurt](#), [Jonathan Fay](#), Erin Johnson, [Susanna Kohler](#), [Knut Olsen](#), [Fred Rasio](#), Doug Roberts, [Philip Rosenfield](#), [Gretchen Stahlman](#), Mark SubbaRao, [Julie Steffen](#), [Matt Turk](#), Patricia Udomprasert, Jaap Vreeling, [A. David Weigel](#), Ryan Wyatt, Martin Woodward, and Curtis Wong. We also thank the the current and past AAS WWT Steering Committee members for support, ideas, and feedback on development initiatives: Amanda Bauer, [Nancy Brickhouse](#), [Jack Burns](#), Buell Jannuzi, Jessica Kirkpatrick, [Sarah Loebman](#), [Kevin Marvel](#), [Megan Schwamb](#), and [Chick](#)

<sup>29</sup> <https://github.com/WorldWideTelescope/pywwt-notebooks>

Woodward. We thank Doug Roberts and Julie Steffen for AAS WWT project leadership.

Chen-Zhou Cui is thankful for support from the National Natural Science Foundation of China (NSFC)(11503051, U1531111, U1531115, U1531246, U1731125, U1731243). Chen-Zhou Cui also thanks the National R&D Infrastructure and Facility Development Program of China, “Earth System Science Data Sharing Platform” and “Fundamental Science Data Sharing Platform” (DKA2017-12-02-XX).

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

We acknowledge the use of the Legacy Archive for Microwave Background Data Analysis (LAMBDA), part of the High Energy Astrophysics Science Archive Center (HEASARC). HEASARC/LAMBDA is a service of the Astrophysics Science Division at the NASA Goddard Space Flight Center.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS website is <http://www.sdss.org/>. The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

The Digitized Sky Survey was produced at the Space Telescope Science Institute under U.S. Government grant NAG W-2166. The images of these surveys are based on photographic data obtained using the Oschin Schmidt Telescope on Palomar Mountain and the UK Schmidt Telescope. The plates

were processed into the present compressed digital form with the permission of these institutions.

This research has made use of the USNO Image and Catalog Archive operated by the United States Naval Observatory, Flagstaff Station (<http://www.nofs.navy.mil/data/fchpix/>) from Monet (1998).

This work is based on observations made with the NASA *Galaxy Evolution Explorer*. *GALEX* is operated for NASA by the California Institute of Technology under NASA contract NAS5-98034.

Some of the data presented in this paper were obtained from the Mikulski Archive for Space Telescopes (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-*HST* data is provided by the NASA Office of Space Science via grant NNX13AC07G and by other grants and contracts.

The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

This research has made use of data obtained from or software provided by the US National Virtual Observatory, which is sponsored by the National Science Foundation.

We acknowledge the use of public data from the *Swift* data archive.

This research has made use of data and/or services provided by the International Astronomical Union’s Minor Planet Center.

This research uses services or data provided by the Science Data Archive at NOAO. NOAO is operated by the Association of Universities for Research in Astronomy (AURA), Inc. under a cooperative agreement with the National Science Foundation.

This work made use of the IPython package (Pérez & Granger 2007), Scikit-learn (McKinney 2010), and Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013).

The acknowledgments were (partially) compiled using the Astronomy Acknowledgment Generator.

*Facilities:* *CGRO*, *COBE*, *EUVE*, *Fermi*, *GALEX*, *GRANAT*, HEASARC, IRSA, MAST, NRAO, OCA:*IRIS*, *Planck*, *ROSAT*, Sloan, *Swift*, VLA, *WISE*, *WMAP*, WSRT.

*Software:* Astropy (Astropy Collaboration et al. 2013); pyWWT (Robitaille et al. 2017).

## Appendix Curated Data

Below are tables of data curated by WorldWide Telescope at the time of publication (as opposed to data WorldWide Telescope accesses over the Internet).

**Table 4**  
All-Sky Surveys

Name (Bandpass)	Approximate TOAST Resolution (arcsec pixel <sup>-1</sup> )	Notes, Credits
Radio		
VLSS: VLA Low-frequency Sky Survey	79.2	Cohen et al. (2007) <sup>a</sup>
VLA FIRST: Faint Images of the Radio Sky at Twenty-centimeters	9.90	Becker et al. (1995)
NVSS: NRAO VLA Sky Survey	19.8	Condon et al. (1998) <sup>a</sup>
SUMSS: Sydney University Molonglo Sky Survey	19.8	Bock et al. (1999) (SUMSS, Min = 0.001, Max = 10). <sup>a</sup>
Westerbork Northern Sky Survey	9.90	WENSS Team. The WENSS project is a collaboration between the Netherlands Foundation for Research in Astronomy (NFRA/ASTRON) and the Leiden Observatory. (WENSS, Min = 0.001, Max = 10), who got it from the WENSS FTP site 1999 Mar 18. The original data were found using the Westerbork Synthesis Radio Telescope, which is operated by NFRA/ASTRON with support from the Netherlands Foundation for Scientific Research (NWO). <sup>a</sup>
Bonn 1420 MHz Survey	39.6	Max Planck Institute for Radio Astronomy, generated by Patricia Reich and Wolfgang Reich using the Bonn Stockert 25 m telescope. (1420 MHz, Min = 3, Max = 35). <sup>a</sup>
HI All-Sky Continuum Survey	158	Max Planck Institute for Radio Astronomy, generated by Glyn Haslam using data taken at the Jodrell Bank, Effelsberg and Parkes telescopes. (408 MHz, Min = 10, Max = 200). <sup>a</sup>
Microwave		
<i>Planck</i>	39.6	<i>Planck</i> is a European Space Agency mission, with significant participation from NASA. NASA's <i>Planck</i> Project Office is based at JPL. JPL contributed mission-enabling technology for both of <i>Planck</i> 's science instruments. European, Canadian and U.S. <i>Planck</i> scientists work together to analyze the <i>Planck</i> data.
CMB		
Dust and Gas		
Thermal Dust		
Spinning Dust		
Molecular Gas (CO)		
Ionized Gas (free-free)		
Synchrotron (non-thermal)		
Lensing (Mass)		
<i>WMAP</i>	158	NASA/ <i>WMAP</i> Science Team
ILC 5 year Cosmic Microwave Background		
K Band, Linear, Nonlinear (23 GHz), and Polarization Map (50 uK)		
Ka Band, Linear, Nonlinear (33 GHz), and Polarization Map (35 uK)		
Q Band, Linear, Nonlinear (41 GHz), and Polarization Map (35 uK)		
V Band, Linear, Nonlinear (61 GHz), and Polarization Map (35 uK)		
W Band, Linear, Nonlinear (94 GHz), and Polarization Map (35 uK)		
<i>WMAP</i> QVW (Power and Linear)		
Infrared		
SFD Dust Map, 100 Micron	39.6	Data provided by two NASA satellites, the <i>Infrared Astronomy Satellite (IRAS)</i> and the <i>Cosmic Background Explorer (COBE)</i> . Processing was performed by David J. Schlegel, Douglas P. Finkbeiner and Marc Davis, at Princeton University and University of California, Berkeley. (SFD100m, Min = 5, Max = 5). <sup>a</sup>
<i>WISE</i> All Sky	19.8	NASA/JPL-Caltech/UCLA
2MASS Two Micron All Sky Survey		Synthetic catalog image by Jina Suh from 2MASS catalog.
Imagery	9.90	
Synthetic catalog	39.6	
<i>IRIS</i> : Improved Reprocessing of <i>IRAS</i> Survey, 12, 25, 60, and 100 $\mu$ m	39.6	Red is IRIS100, green is IRIS60, blue is IRIS12 (IRIS12: Min = 0.5, Max = 1; IRIS25: Min = 1, Max = 2; IRIS60: Min = 1, Max = 2; IRIS100: Min = 4, Max = 4). <sup>a,b</sup>
<i>COBE</i> Diffuse Infrared Background Experiment (DIRBE)	39.6	NASA/ <i>COBE</i> ( <i>COBE</i> , Min = 10, Max = 5). <sup>a</sup>
<i>COBE</i> DIRBE Annual Average Map	39.6	NASA/ <i>COBE</i> (COBEAAM, Min = 10, Max = 5). <sup>a</sup>
<i>COBE</i> DIRBE Zodi-Subtracted Mission Average	39.6	NASA/ <i>COBE</i> (COBEZSMA, Min = 10, Max = 5). <sup>a</sup>



**Table 4**  
(Continued)

Name (Bandpass)	Approximate TOAST Resolution (arcsec pixel <sup>-1</sup> )	Notes, Credits
Optical		
Hydrogen Alpha Full Sky Map	39.6	The full-sky H-alpha map (6' FWHM resolution) is a composite of the Virginia Tech Spectral line Survey (VTSS) in the North and the South H-Alpha Sky Survey Atlas (SHASSA). The Wisconsin H-Alpha Mapper (WHAM) survey provides a stable zero-point over three-fourths of the sky on a one-degree scale.
SDSS: Sloan Digital Sky Survey Digitized Sky Survey	0.30	
Tycho (Synthetic)	79.2	NASA/Goddard Space Flight Center Scientific Visualization Studio.
USNOB: US Naval Observatory B 1.0 (Synthetic)	39.6	Monet (1998).
Ultraviolet		
<i>GALEX</i> 2 Combined	2.48	GR2/3 release <sup>a,c</sup>
<i>GALEX</i> 2 Near-UV	2.48	GR2/3 release <sup>a,c</sup>
<i>GALEX</i> 2 Far-UV	2.48	GR2/3 release <sup>a,c</sup>
<i>GALEX</i> 4 Near-UV	2.48	Near-UV band (1770–2730 Å) of the GR4 release <sup>a,c</sup>
<i>GALEX</i> 4 Far-UV	2.48	Far-UV band (1350–1780 Å) of the GR4 release <sup>a,c</sup>
<i>Extreme Ultraviolet Explorer (EUVE)</i>	79.2	Red is 555 Å (euve555), green is 405 Å, and red is 83 Å. <sup>a,c,d</sup>
<i>EUVE</i> : 83 Å	79.2	(euve83, Min = 1, Max = 150). <sup>a,c,d</sup>
<i>EUVE</i> : 171 Å	79.2	(euve171, Min = 4, Max = 200). <sup>a,c,d</sup>
<i>EUVE</i> : 405 Å	79.2	(euve405, Min = 0.7, Max = 70). <sup>a,c,d</sup>
<i>EUVE</i> : 555 Å	79.2	(euve555, Min = 5, Max = 500). <sup>a,c,d</sup>
X-Ray		
RASS: <i>ROSAT</i> All Sky Survey	19.8	This is a composite of three RASS3 surveys from the <i>ROSAT</i> Data Archive of the Max-Planck-Institut für extraterrestrische Physik (MPE) at Garching, Germany. Red is the soft band (RASS3sb), green is the broad band (RASS3bb), blue is the hard band (RASS3hb) <sup>a</sup>
<i>ROSAT</i> Hard Band Count Map	19.8	(RASS3hb, Min = 0.2, Max = 50). <sup>a,c</sup>
<i>ROSAT</i> Soft Band Count Map	19.8	(RASS3sb, Min = 0.2, Max = 50). <sup>a,c</sup>
<i>ROSAT</i> Broad Band Count Map	19.8	(RASS3bb, Min = 0.2, Max = 50). <sup>a,c</sup>
<i>ROSAT</i> Soft Band Intensity	19.8	(RASS3sb, Min = 0.2, Max = 10). <sup>a,c</sup>
<i>ROSAT</i> Hard Band Intensity	19.8	(RASS3hb, Min = 0.2, Max = 10). <sup>a,c</sup>
<i>ROSAT</i> PSPC summed pointed observations, 2° cutoff, intensity	9.90	Observational data from NASA Goddard Space Flight Center, (PSPC2int, Min = 0.02, Max = 0.01). <sup>a</sup>
<i>Swift</i> BAT All-Sky Survey: Significance 14– 195 keV	79.2	NASA BAT Team. (BATSig, Min = 1, Max = 10). <sup>a</sup>
<i>Swift</i> BAT All-Sky Survey: Flux 14–195 keV	79.2	NASA BAT Team. (BATFlux, Min = 1e–5, Max = 1e–3). <sup>a</sup>
<i>GRANAT</i> /SIGMA Significance	79.2	(GRANAT_SIGMA_sig, Min = 0.5, Max = 50). <sup>a,f</sup>
<i>GRANAT</i> /SIGMA Flux	79.2	(GRANAT_SIGMA_flux, Min = 1e–5, Max = 0.01). <sup>a,f</sup>
Gamma-Ray		
<i>Fermi</i>	158	NASA and the <i>Fermi</i> -LAT Team.
<i>Fermi</i> Six Months	158	NASA and the <i>Fermi</i> -LAT Team.
<i>Fermi</i> Year Two	39.6	
<i>Fermi</i> Year Three	79.2	
<i>Fermi</i> LAT Year Eight	39.6	
<i>CGRO</i> Compton Telescope: 3 channel data	79.2	CompTel Instrument Team. Maps generated by Andrew Strong, Max-Planck Institute for Extraterrestrial Physics, Garching, Germany. (comptel, Min = 0.05, Max = 0.1). <sup>a</sup>
EGRET Soft	79.2	EGRET Instrument team, NASA Goddard Space Flight Center. (EGRET soft, Min = 0.1, Max = 0.002). <sup>a</sup>
EGRET Hard	79.2	EGRET Instrument team, NASA Goddard Space Flight Center. (EGRET hard, Min = 0.02, Max = 0.005). <sup>a</sup>

**Notes.**<sup>a</sup> TOAST-formatted mosaics were obtained using facilities of NASA's SkyView Virtual Telescope.<sup>b</sup> Original *IRAS* data: NASA/JPL IPAC; *IRIS* Reprocessing: Canadian Institute for Theoretical Astrophysics/Institut d'Astrophysique Spatiale.<sup>c</sup> Data archived at MAST/STScI.<sup>d</sup> Center for Extreme UV Astronomy, University of California at Berkeley.<sup>e</sup> *ROSAT* Data Archive of the Max-Planck-Institut für extraterrestrische Physik (MPE) at Garching, Germany.<sup>f</sup> High Energy Astrophysics Department, Space Research Institute, Moscow, Russia; CEA, Centre d'Etudes de Saclay Orme des Merisiers, France; Centre d'Etude Spatiale des Rayonnements, Toulouse, France; Federation de Recherche Astroparticule et Cosmologie Universite de Paris, France.

**Table 5**  
Panoramas

Name	Approximate TOAST Resolution (arcsec pixel <sup>-1</sup> )	Notes, Credits
Apollo 12: Landing Site	19.8	Panorama taken by Apollo 12 Lunar Module Pilot Alan Bean in 1969 and assembled at the NASA Ames Research Center in 2007.
Apollo 17: Shorty Crater	19.8	Panorama taken by Apollo 17 Commander Gene Cernan in 1972 and assembled at the NASA Ames Research Center in 2007.
Pathfinder: Improved MPF 360° Presidential Panorama	79.1	NASA/JPL. This is the 1993 “geometrically improved, color enhanced” version of the 360° “Gallery Pan” at Ares Vallis, the first contiguous, uniform panorama taken by the Imager for Mars (IMP) over the course of Sols 8, 9, and 10.
Pathfinder: “Many Rovers”	79.1	Panoramas made available by Dr. Carol Stoker, NASA AMES. NASA/JPL/USGS.
Pathfinder: Monster (stereo)	79.1	
Pathfinder: Landing site from Sagan Memorial Station (stereo)	39.6	
<i>Spirit</i> : McMurdo	19.8	This 360° view, called the “McMurdo” panorama, comes from the panoramic camera (Pancam) on NASA’s Mars Exploration Rover <i>Spirit</i> . From April through 2006 Oct, <i>Spirit</i> has stayed on a small hill known as “Low Ridge.” There, the rover’s solar panels are tilted toward the Sun to maintain enough solar power for <i>Spirit</i> to keep making scientific observations throughout the winter on southern Mars. This view of the surroundings from <i>Spirit</i> ’s “Winter Haven” is presented in approximately true color. Image mosaicking: Kris Kapraro, Bob Deen, and the JPL/MIPL team. <sup>a,b</sup>
<i>Spirit</i> : McMurdo (false color)	19.8	The Pancam began shooting component images of this panorama during <i>Spirit</i> ’s sol 814 (2006 Apr 18) and completed the part shown here on sol 932 (2006 Aug 17). The panorama was acquired using all 13 of the Pancam’s color filters. Image mosaicking: Kris Kapraro, Bob Deen, and the JPL/MIPL team. <sup>a,b</sup>
<i>Spirit</i> : McMurdo (color stereo)	19.8	This 360° view comes from the panoramic camera (Pancam) on NASA’s Mars Exploration Rover <i>Spirit</i> . From 2006 Apr through Oct, <i>Spirit</i> has stayed on a small hill known as “Low Ridge.”
<i>Spirit</i> : McMurdo (left eye)	19.8	The Pancam on the <i>Spirit</i> Rover began shooting component images of this panorama during <i>Spirit</i> ’s sol 814 (2006 Apr 18) and completed the part shown here on sol 932 (2006 Aug 17). The panorama was acquired using all 13 of the Pancam’s color filters, using lossless compression for the red and blue stereo filters, and only modest levels of compression on the remaining filters. The overall panorama consists of 1449 Pancam images. <sup>a</sup>
<i>Spirit</i> : McMurdo (right eye)	19.8	NASA’S Mars Exploration Rover <i>Spirit</i> captured this westward view from atop a low plateau where <i>Spirit</i> spent the closing months of 2007.
<i>Spirit</i> : West Valley	19.8	
<i>Spirit</i> : West Valley (false color)	19.8	<i>Spirit</i> acquired the 405 individual images that make up this 360° view of the surrounding terrain using five different filters on the panoramic camera. The rover took the images on Martian days, or sols, 672–677 (2005 Nov 23–28). <sup>a,b,c</sup>
<i>Spirit</i> : West Valley (stereo)	19.8	
<i>Spirit</i> : Descent from Husband Hill	19.8	
<i>Spirit</i> : Descent from Husband Hill (false color)	19.8	NASA’s Mars Exploration Rover <i>Spirit</i> acquired this high-resolution view of intricately layered exposures of rock while parked on the northwest edge of the bright, semi-circular feature known as “Home Plate.” The rover was perched at a 27° upward tilt while creating the panorama, resulting in the “U” shape of the mosaic. <sup>d</sup>
<i>Spirit</i> : Descent from Husband Hill (stereo)	19.8	
<i>Spirit</i> : Gibson Panorama at Home Plate (false color)	19.8	
<i>Spirit</i> : Gibson Panorama at Home Plate	39.6	This is the <i>Spirit</i> Pancam “Home Plate South” panorama, acquired on sols 1325–1332 (2007 Sep 25–Oct 2). This is an approximate true-color rendering using Pancam’s 753, 535, and 432 nm filters. <sup>a</sup>
<i>Spirit</i> : Home Plate South	19.8	
<i>Spirit</i> : Home Plate South (false color)	19.8	This is a false-color version of the <i>Spirit</i> Pancam “Home Plate South” panorama, acquired on sols 1325–1332 (2007 Sep 25–Oct 2). <sup>a</sup>
<i>Spirit</i> : Home Plate South (stereo)	19.8	The panoramic camera on NASA’s Mars Exploration Rover <i>Spirit</i> took the hundreds of images combined into this 360° view, the “Husband Hill Summit” panorama. The images were acquired on <i>Spirit</i> ’s sols 583–586 (2005 Aug 24–27), shortly after the rover reached the crest of “Husband Hill” inside Mars’ Gusev Crater.
<i>Spirit</i> : Husband Hill Summit	39.6	
<i>Spirit</i> : Santa Anita	19.8	This color mosaic was taken on 2004 May 21, 25, and 26 by the panoramic camera on NASA’s Mars Exploration Rover <i>Spirit</i> and was acquired from a position roughly three-fourths the way between “Bonneville Crater” and the base of the “Columbia Hills.” <sup>a</sup>
<i>Spirit</i> : Independence	19.8	The 108 images used to make this panorama were obtained on sols 536–543 (2005 Jul 6–13) from a position in the Columbia Hills near the summit of Husband Hill.

**Table 5**  
(Continued)

Name	Approximate TOAST Resolution (arcsec pixel <sup>-1</sup> )	Notes, Credits
<i>Spirit</i> : Everest	19.8	It took <i>Spirit</i> three days, sols 620–622 (2005 Oct 1–3), to acquire the 81 images combined into this mosaic, looking outward in every direction from the true summit of Husband Hill. The Sky was fixed by Jim St George. <sup>a,b,c</sup>
<i>Spirit</i> : Everest (stereo)	19.8	Composed of 81 images acquired on sols 620–622 (2005 Oct 1–3) from a position in the Columbia Hills at the true summit of Husband Hill. <sup>a,c</sup>
<i>Spirit</i> Mission Success	39.6	This panorama was acquired in eight parts called octants, over the course of three different sols. Because the octants were taken at different times of day and across different days when the dust abundance was changing, there were large brightness and color seams between the octants in the assembled mosaic. <sup>a</sup>
<i>Spirit</i> : Thanksgiving	19.8	This is the <i>Spirit</i> Pancam “Thanksgiving” panorama, acquired on sols 318–325 (2004 Nov 24 to Dec 2) from a position along the flank of Husband Hill, which is the peak just left of the center of this mosaic, just east of the West Spur of the Columbia Hills. <sup>a,b,c</sup>
<i>Spirit</i> : Thanksgiving (stereo)	79.1	
<i>Spirit</i> : Cahokia	19.8	The approximate true-color image, nicknamed the “Cahokia panorama” after the Native American archaeological site near St. Louis, was acquired between sols 213 to 223 (2004 Aug 9–19). The panorama consists of 470 images acquired through six panoramic camera filters (750–480 nm). Stereo images: JPL/MIPL (Bob Deen). <sup>a</sup>
<i>Spirit</i> : Cahokia (stereo, unadjusted)	19.8	
<i>Spirit</i> : Cahokia (stereo, tilt-adjusted)	158	
<i>Spirit</i> : Larry’s Lookout	19.8	This is the <i>Spirit</i> Pancam “Lookout” panorama, acquired on sols 410–413 (2005 Feb 27 to Mar 2) from a position known informally as “Larry’s Lookout” along the drive up Husband Hill. <sup>a,b,c</sup>
<i>Spirit</i> : Larry’s Lookout (stereo)	19.8	
<i>Spirit</i> : Whale of a Panorama	19.8	<i>Spirit</i> produced this 220° image mosaic two-thirds of the way to the summit of Husband Hill from images collected from sol 497 to 500 (2005 May 27 through May 30). <sup>a</sup>
<i>Spirit</i> : From The Summit	39.6	This approximate true-color 240° panorama was taken by NASA’s <i>Spirit</i> rover from the top of “Husband Hill” in the “Columbia Hills” of Gusev Crater. The mosaic is made up of images taken by the rover’s panoramic camera over a period of three days (sols 583–585, or 2005 Aug 24–26).
<i>Spirit</i> : Paige Panorama	39.6	This 230° panorama was composed from 72 images from the Pancam on <i>Spirit</i> on 2006 Feb 19. <sup>d</sup>
<i>Spirit</i> : Paige Panorama (false color)	39.6	
<i>Spirit</i> : Legacy	19.8	The 78 images used to make this were acquired on sols 59–61 (2004 Mar 3–5) from a position about halfway between the landing site and the rim of Bonneville crater, within the transition from the relatively smooth plains to the more rocky and rugged ejecta blanket of Bonneville. <sup>a,b,c</sup>
<i>Spirit</i> : Bonneville	19.8	The rim and interior of a crater nicknamed “Bonneville” dominate this 180°, false-color mosaic. <i>Spirit</i> recorded this view on the rover’s 68th sol, 2004 Mar 12, one sol after reaching this location. The rover remained here in part to get this very high-resolution, color mosaic, from which scientists can gain insight about the depth of the surface material at Bonneville and make future observation plans. On sol 71, <i>Spirit</i> was instructed to drive approximately 15 m (49 feet) along the crater rim to a new vantage point. <sup>a</sup>
<i>Opportunity</i> : Erebus	19.8	This panorama was made from 635 images acquired on sols 652–663 (2005 Nov 23 to Dec 5), as NASA’s Mars Exploration Rover <i>Opportunity</i> was exploring sand dunes and outcrop rocks in Meridiani Planum. Image mosaicking: JPL/MIPL Team (Bob Deen, Oleg Pariser, Jeffrey Hall). <sup>a,b</sup>
<i>Opportunity</i> : Erebus (Stereo)	19.8	
<i>Opportunity</i> : Lyell	39.6	This view combines many images taken by <i>Opportunity</i> ’s panoramic camera (Pancam) from the 1332nd through 1379th Martian days, or sols, of the mission (2007 Oct 23 to Dec 11). The main body of the crater appears in the upper right of this stereo panorama, with the far side of the crater lying about 800 m (half a mile) away. Bracketing that part of the view are two promontories on the crater’s rim at either side of Duck Bay. They are “Cape Verde,” about 6 m (20 feet) tall, on the left, and “Cabo Frio,” about 15 m (50 feet) tall, on the right. The rest of the image, other than the sky and portions of the rover, is ground within Duck Bay. Image mosaicking: Jim Bell and Jonathan Joseph. <sup>a,b</sup>
<i>Opportunity</i> : Lyell (Stereo)	19.8	
<i>Opportunity</i> : Rub al Khali	19.8	Taken on the plains of Meridiani during the period from the rover’s 456th to 464th sols on Mars (2005 May 6–14). <i>Opportunity</i> was about 2 km (1.2 miles) south of “Endurance Crater” at a place known informally as “Purgatory Dune.” <sup>a</sup>
<i>Opportunity</i> : Endurance South	19.8	This 360° panorama shows “Endurance Crater” and the surrounding plains of Meridiani Planum. This is the second large panoramic camera mosaic of Endurance, and was obtained from a high point near the crater’s south rim. <sup>a</sup>
<i>Opportunity</i> : Endurance South (false color)	19.8	



**Table 5**  
(Continued)

Name	Approximate TOAST Resolution (arcsec pixel <sup>-1</sup> )	Notes, Credits
<i>Opportunity</i> Mission Success	19.8	This expansive view of the Martian real estate surrounding the Mars Exploration Rover <i>Opportunity</i> is the first 360°, high-resolution color image taken by the rover's panoramic camera. The airbag marks, or footprints, seen in the soil trace the route by which <i>Opportunity</i> rolled to its final resting spot inside a small crater at Meridiani Planum, Mars. The exposed rock outcropping is a future target for further examination. This image mosaic consists of 225 individual frames.
<i>Opportunity</i> : Burns Cliff	39.6	This is a view of "Burns Cliff" after driving right to the base of this southeastern portion of the inner wall of "Endurance Crater." The view combines 46 images taken by <i>Opportunity</i> 's panoramic camera between the rover's 287th and 294th Martian days (2004 Nov 13–20). <sup>a</sup>
<i>Opportunity</i> : Payson Panorama	79.1	The panoramic camera on board NASA's Mars Exploration Rover <i>Opportunity</i> acquired this panorama of the Payson outcrop on the western edge of Erebus Crater during <i>Opportunity</i> 's sol 744 (2006 Feb 26). USGS. <sup>a</sup>
<i>Opportunity</i> : Beagle Crater	19.8	<i>Opportunity</i> took the mosaic of images that make up this 360° view of the rover's surroundings with the panoramic camera on the rover's 901st through 904th sols, or Martian days (2006 Aug 6 through Aug 9), of exploration. This is an approximate true-color image combining exposures taken through the panoramic camera's 753 nm, 535 nm, and 432 nm filters. Image mosaicking: Cornell Pancam team. NASA/JPL/Cornell/UNM. <sup>b</sup>
<i>Opportunity</i> : Beagle Crater (false color)	19.8	
<i>Opportunity</i> : Lion King	19.8	This shows "Eagle Crater" and the surrounding plains of Meridiani Planum. It was stitched from 558 images obtained on sols 58 and 60 by the Mars Exploration Rover <i>Opportunity</i> 's panoramic camera. <sup>a</sup>
Phoenix: Landing Site	39.6	This view combines more than 400 images taken during the first several weeks after NASA's Phoenix Mars Lander arrived on an arctic plain at 68°22 north latitude, 234°25 east longitude on Mars.









**Notes.**<sup>a</sup> NASA/JPL/Cornell.<sup>b</sup> Calibration and color rendering: Cornell Calibration Crew and the Pancam team (Jim Bell).<sup>c</sup> Image mosaicking: JPL/MIPL (Bob Deen).<sup>d</sup> NASA/JPL-Caltech/USGS/Cornell.**Table 6**  
Planet Maps

Name	Approximate TOAST Resolution (arcsec pixel <sup>-1</sup> )	Notes, Credits
Bing Maps Aerial	0.002	
Bing Maps Hybrid	0.002	
Bing Maps Streets	0.002	
Earth at Night 2012	4.94	NASA Earth Observatory/NOAA NGDC.
Earth at Night	19.8	Data are courtesy of Marc Imhoff of NASA GSFC and Christopher Elvidge of NOAA NGDC. Image by Craig Mayhew and Robert Simmon, NASA GSFC.
Blue Marble 2004 Jan	19.8	Blue Marble Next Generation, 2004 Jan. Visible Earth Project, NASA.
Blue Marble 2004 Jul	19.8	Blue Marble Next Generation, 2004 Jul. Visible Earth Project, NASA.
Open Street Maps	0.010	OpenStreetMap contributors, CC BY-SA.
Open Street Maps Bike Map	0.010	OpenStreetMap contributors, CC BY-SA.
Moon	2.47	Image courtesy of NASA/JPL.
Mercury	9.89	NASA/Johns Hopkins University Applied Physics Laboratory/Carnegie Institution of Washington.
Venus	79.1	Image courtesy of NASA/JPL/Space Science Institute.
Mars Visible Imagery	9.89	NASA/USGS/Malin Space Science Systems/JPL.
Jupiter	158	Image courtesy of NASA/JPL/Space Science Institute.
Phobos (Mars)	316	Viking images of the moon Phobos. <sup>a</sup>
Deimos (Mars)	316	Viking images of the moon Deimos. <sup>a</sup>
Io (Jupiter)	79.1	Images courtesy of NASA/JPL/Space Science Institute.
Europa (Jupiter)	79.1	
Ganymede (Jupiter)	79.1	

**Table 6**  
(Continued)

Name	Approximate TOAST Resolution (arcsec pixel <sup>-1</sup> )	Notes, Credits
Callisto (Jupiter)	79.1	
Mimas (Saturn)	316	Approximately half the surface of moon Mimas. <sup>a,b</sup>
Enceladus (Saturn)	316	Approximately half the surface of moon Enceladus. <sup>a,b</sup>
Tethys (Saturn)	316	Most of the surface of the moon Tethys. <sup>a,b</sup>
Dione (Saturn)	316	Most of the surface of the moon Dione. <sup>a,b</sup>
Rhea (Saturn)	316	The moon Rhea. <sup>a,b</sup>
Iapetus (Saturn)	316	Approximately half of the surface of the moon Iapetus. <sup>a,b</sup>
Ariel (Uranus)	316	Most of the Southern Hemisphere of the moon Ariel. <sup>a,b</sup>
Umbriel (Uranus)	316	Most of the Southern Hemisphere of the moon Umbriel. <sup>a,b</sup>
Titania (Uranus)	316	Most of the Southern Hemisphere of the moon Titania. <sup>a,b</sup>
Oberon (Uranus)	316	Most of the Southern Hemisphere of the moon Oberon. <sup>a,b</sup>
Miranda (Uranus)	316	Most of the Southern Hemisphere of the moon Miranda. <sup>a,b</sup>

**Notes.**<sup>a</sup> Caltech/JPL/USGS.<sup>b</sup> From *Voyager* mission photographs.**ORCID iDs**

Philip Rosenfield  <https://orcid.org/0000-0001-9306-6049>  
Jonathan Fay  <https://orcid.org/0000-0003-2500-8984>  
Ronald K Gilchrist  <https://orcid.org/0000-0001-8246-5001>  
Chenzhou Cui  <https://orcid.org/0000-0002-7456-1826>  
A. David Weigel  <https://orcid.org/0000-0002-8026-2291>  
Thomas Robitaille  <https://orcid.org/0000-0002-8642-1329>  
Oderah Justin Otor  <https://orcid.org/0000-0002-4679-5692>  
Alyssa Goodman  <https://orcid.org/0000-0003-1312-0477>

**References**

- Acton, C., Bachman, N., Semenov, B., & Wright, E. 2017, *P&SS*, **150**, 9  
Acton, C. H. 1996, *P&SS*, **44**, 65  
Agarwal, D., Cheah, Y.-W., Fay, D., et al. 2011, *The International Journal of High Performance Computing Applications*, **25**, 304  
Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, **558**, A33  
Batygin, K., & Brown, M. E. 2016, *AJ*, **151**, 22  
Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, **450**, 559  
Berriman, G. B., & Good, J. C. 2017, *PASP*, **129**, 058006  
Bock, D. C.-J., Large, M. I., & Sadler, E. M. 1999, *AJ*, **117**, 1578  
Cohen, A. S., Lane, W. M., Cotton, W. D., et al. 2007, *AJ*, **134**, 1245  
Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, *AJ*, **115**, 1693  
Cui, C., & Li, S. 2018, arXiv:1801.05098  
Currie, T., Lisse, C. M., Kuchner, M., et al. 2015, *ApJL*, **807**, L7  
Geller, M. J., & Huchra, J. P. 1989, *Sci*, **246**, 897  
Gray, J., & Szalay, A. S. 2004, arXiv:cs/0403018  
Jurić, M., Kantor, J., Lim, K., et al. 2015, arXiv:1512.07914  
Kluyver, T., Ragan-Kelley, B., Pérez, F., et al. 2016, in *Positioning and Power in Academic Publishing: Platters, Agents and Agendas*, ed. F. Loizides & B. Schmidt (Amsterdam: IOS Press), 87  
Kunszt, P. Z., Szalay, A. S., Csabai, I., & Thakar, A. R. 2000, in *ASP Conf. Ser. 216, Astronomical Data Analysis Software and Systems IX*, ed. N. Manset, C. Veillet, & D. Crabtree (San Francisco, CA: ASP), 141  
Ladd, E. F., Gingrich, E. C., Nottis, K. E. K., Udomprasert, P., & Goodman, A. A. 2015, in *ASP Conf. Ser. 500, Celebrating Science: Putting Education Best Practices to Work*, ed. G. Shultz et al. (San Francisco, CA: ASP), 191  
McKinney, W. 2010, in *Proc. 9th Python in Science Conf.*, ed. S. van der Walt & J. Millman, 51  
Monet, D. 1998, *USNO-A2.0* (Washington, DC: United States Naval Observatory) (<http://tdc-www.harvard.edu/catalogs/ua2.html>)  
Nakata, M., Hamacher, D. W., Warren, J., et al. 2014, *Australian Academic & Research Libraries*, **45**, 101  
Pérez, F., & Granger, B. E. 2007, *CSE*, **9**, 21  
Perryman, M. A. C., Lindegren, L., Kovalevsky, J., et al. 1997, *A&A*, **323**, L49  
Qiao, C., Cui, C., Zheng, X., & Xu, Y. 2011, in *Proc. of the 5th Int. Technology, Education and Development Conf.*, ed. L. Gómez Chova, I. Candel Torres, & A. López Martínez (Valencia, Spain: IATED), 128, (<https://library.iated.org/view/QIAO2011THE>)  
Roberts, M. J. 2014, in *ASP Conf. Ser. 483, Ensuring Stem Literacy: A National Conference on STEM Education and Public Outreach*, ed. J. G. Manning et al. (San Francisco, CA: ASP), 223  
Robitaille, T., Otor, O. J., ZuHone, J., & Rosenfield, P. 2017, *PyWWT: WorldWide Telescope from Python*, Zenodo, doi:10.5281/zenodo.1155830  
Rosenfield, P., Gaily, J., Fraser, O., & Wisniewski, J. 2014, *CAPJ*, **15**, 35  
Sanders, N. E., Faesi, C., & Goodman, A. A. 2014, *JSEdT*, **23**, 431  
Scoville, N., Aussel, H., Brusa, M., et al. 2007, *ApJS*, **172**, 1  
Udomprasert, P., Goodman, A., Sunbury, S., et al. 2014, in *ASP Conf. Ser. 483, Ensuring Stem Literacy: A National Conference on STEM Education and Public Outreach*, ed. J. G. Manning et al. (San Francisco, CA: ASP), 297  
Udomprasert, P. S., Goodman, A. A., Plummer, J., et al. 2016, in *American Astronomical Society Meeting Abstracts 227*, 229.01  
Udomprasert, P. S., Goodman, A. A., & Wong, C. 2012, in *ASP Conf. Ser. 457, Connecting People to Science: A National Conference on Science Education and Public Outreach*, ed. J. B. Jensen et al. (San Francisco, CA: ASP), 149  
Wang, Q., Qiao, C., & Zheng, X. 2015, *International Journal of Learning and Teaching*, **1**, 134  
Wong, C. 2008, in *ASP Conf. Ser. 389, EPO and a Changing World: Creating Linkages and Expanding Partnerships*, ed. C. Garmany, M. G. Gibbs, & J. W. Moody (San Francisco, CA: ASP), 227  
Yusef-Zadeh, F., Roberts, D. A., Wardle, M., et al. 2015, *ApJL*, **801**, L26